



HOME | SUBMIT | FAQ | BLOG | ALERTS / RSS | ABOUT | CHANNELS

Search Q

Advanced Search

New Results

▲ Follow this preprint

Bile Acids Serve As Endogenous Antagonists Of The Leukemia Inhibitory Factor (LIF) Receptor in Oncogenesis

© Cristina Di Giorgio, Elva Morretta, Antonio Lupia, Rachele Bellini, Carmen Massa, Ginevra Urbani, Martina Bordoni, Silvia Marchianò, Pasquale Rapacciuolo, Claudia Finamore, Valentina Sepe, Maria Chiara Monti, Federica Moraca, Nicola Natalizi, Luigina Graziosi, Eleonora Distrutti, Michele Biagioli, Bruno Catalanotti, Annibale Donini, Angela Zampella, Stefano Fiorucci

doi: https://doi.org/10.1101/2023.09.30.560269

This article is a preprint and has not been certified by peer review [what does this mean?].



Abstract

Full Text

Info/History

Metrics

Preview PDF

Summary

The leukemia inhibitory factor (LIF) is member of IL-6 family of cytokines involved immune regulation, morphogenesis and oncogenesis. In cancer tissues, LIF binds a heterodimeric receptor (LIFR), formed by a LIFR β subunit and glycoprotein(gp)130, promoting epithelial mesenchymal transition and cell growth. Bile acids are



Citation Tools









COVID-19 SARS-CoV-2 preprints from medRxiv and bioRxiv

Subject Area

Cancer Biology

Subject Areas

All Articles